

References

- FOSTER, J. B. (1965): Mortality and aging of black rhinoceros in East Tsavo Park, Kenya. *E. Afr. Wildlife Journal*, 3, 118—9.
- MERTENS, R. (1966): Zur Typentomologie und Nomenklatur einiger Nashörner der Gattung *Diceros*. *Zool. Garten (N. F.)* 32, 3, 116—7.
- ZUKOWSKY, L. (1964): Die Systematik der Gattung *Diceros* Gray, 1821. *Zool. Garten (N. F.)* 30, 1—2, 1—178.

Author's address: COLIN P. GROVES, Anthropology Department, University of California, Berkeley, Calif. 94720, USA.

Sonderdruck aus *Z. f. Säugetierkunde* Bd. 32 (1967) H. 5, S. 267—276

Mitte Rechte, auch die der Übersetzung, des Nachdrucks und der photomechanischen Wiedergabe, vorbehalten.
 VERLAG PAUL PAREY · HAMBURG 1 · SPIITÄLERSTRASSE 12
 © 1967 Verlag Paul Parey, Hamburg und Berlin

Geographic Variation in the Black Rhinoceros

Diceros bicornis (L., 1758)

By COLIN P. GROVES

Eingang des Ms. 6. I. 1967

The systematics of the Rhinoceroses have been badly neglected in the present century as a whole. There has been no revision of the family Rhinocerotidae, although the interrelationships of living and fossil forms are in constant dispute. Nor has there been much intragenetic taxonomy; only in the past few years has interest seemed to revive in the group. The present author (in press) has attempted a revision of the South-east Asian species *Rhinoceros sondaicus* and *Dicernoceros sumatrensis*, although with the rather small amount of material available this must of necessity be tentative. The only really major work has been that of Zukowsky (1964) on the African species *Diceros bicornis*, the Black Rhinoceros. This has certain drawbacks, but these concern interpretation rather than fact, and as a reference work on aspects of nomenclature, observed variability, distribution etc. it will stand as the source-book on the species for years to come.

Having said this, however, one must proceed to criticise the taxonomic philosophy involved. From a wealth of photographs of living animals and skulls, and from the measurements of 79 skulls, the author felt able to recognise no fewer than sixteen subspecies, as well as a "variety", of which eight were described for the first time. Taking the skull measurements — undeniably the most important line of evidence, if only because subjectivity is absolutely ruled out — this gives an average of just under five skulls per subspecies. When closely examined, the skull measurements do not discriminate between the different forms in many cases, and one is left dependant upon the much more subjective assessment of the photographs to substantiate the forms described.

MERTENS (1966) has already pointed out that three of the names given by Zukowsky have no nomenclatorial status as they were proposed by implication and thus invalid according to the International Code of Zoological Nomenclature, XV congress. Two of these names were based on one skull each; and the third was based on two photographs of freshly-shot specimens. One must always regret the erection of new races on such insecure evidence.

It is for two reasons that the present study is offered: first, for the reason that the taxonomic ideas of Zukowsky will not find ready acceptance among the majority of modern workers; and secondly because a number of specimens not seen by Zukowsky have been studied by the present author. These specimens are from the Powell-Cotton Museum (Birmingham, Kent, England), the Copenhagen Zoological Museum, the Royal College of Surgeons, the Cambridge Zoology Museum, the American Museum of Natural History and the Smithsonian Institution (U.S. National Museum). Also, many of the specimens in Zukowsky's list have been seen by the present author, and the technique of measurement checked; and some additional specimens in London, Munich and Vienna have been seen which were not mentioned in the list. (The London specimens in his cable are from measurements by Hopwood and Schwarz, and the Munich and Vienna ones by Trautner; in the first case, the museum has received

Table 1

Means and standard deviations for population groupings of *Diceros bicornis*. Means only are given where the figure is based on under four specimens; figures in parenthesis indicate that only one specimen could be measured.

	Greatest length	Basal length	Zygomatic breadth	Tarshrow length	Occiput breadth
<i>1. East African area</i>					
Karagwe (4)	586.7 ± 22.5	514.7 ± 8.3	329.8 ± 11.6	252.5 ± 10.7	181.8 ± 16.3
Serengeti (3)	566.7	546.3	329.0	263.5	183.0
Eyasi area (12)	557.2 ± 22.7	525.8 ± 14.8	327.0 ± 12.8	256.9 ± 9.0	187.7 ± 12.6
Arusha (3)	570.5	547.0	333.0	254.0	201.7
Kilimanjaro (3)	559.5	518.5	332.7	245.3	191.3
Tsavu (6)	542.8 ± 15.7	527.5 ± 15.3	332.2 ± 9.0	264.0 ± 13.8	190.5 ± 4.9
Kiboko (4)	546.3 ± 14.9	521.0 ± 5.8	326.8 ± 7.9	244.5 ± 8.8	187.3 ± 9.8
Tana river (9)	523.3	507.7	319.7	243.0	179.0
Lakimunda R. (3)	557.0	534.3	327.3	253.3	183.0
Aberdare (4)	567.5 ± 12.6	546.7 ± 6.2	335.8 ± 6.8	268.7 ± 4.5	179.5 ± 4.3
	594.5	(542.0)	347.5	256.0	206.0
N. Guso Nyiro (6)	578.8 ± 16.5	550.3 ± 9.7	334.8 ± 10.4	249.5 ± 11.3	198.2 ± 19.4
S. Guso Nyiro (3)	557.7	553.3	328.7	256.0	201.0
<i>2. Other areas</i>					
Cape (5)	667.0 ± 37.7	606.3 ± 25.3	372.5 ± 13.4	315.3 ± 17.6	218.8 ± 23.6
S. E. Africa (6)	575.2 ± 8.1	541.7 ± 13.6	350.4 ± 8.5	259.2 ± 11.3	191.3 ± 5.8
Chobe (4)	615.0 ± 15.6	560.0 ± 8.4	352.0 ± 9.9	282.3 ± 7.4	197.5 ± 6.3
Zambesi (8)	567.4 ± 21.8	553.8 ± 20.4	331.1 ± 17.2	249.9 ± 14.4	186.6 ± 12.8
W. Africa (4)	541.3 ± 17.5	522.7 ± 10.8	324.3 ± 3.8	257.5 ± 3.5	191.0 ± 5.5

additional specimens since the two authors wrote, while in the second case it is likely that only the most important skulls were measured).

With the additional material, a more detailed study becomes possible of the populations of East Africa. Table 1 shows the parameters of thirteen East African "populations", while Fig. 1 indicates their geographical placement. (In some cases, individual skulls were available from other localities; these are marked on the map but not given in the table). The first noticeable thing is that there is a clinal size decrease from west to east, from Karagwe to the Tana river. The two skulls from near Lake Baringo are the largest of all, and the Aberdare and Lakimunda populations are also intermediate. There is much less variation in Basal length than in the Greatest length (Occipitonasal), while fluctuations in the other three measurements seem to be more random. Certain populations stand out in one or two features: the Baringo skulls in their broad zygomata and occiput, and the Tana river ones in their narrowness. The five measurements were the most significant of those from ZUKOWSKY'S table; most of the others were only given in a few cases, but some, such as occipital height, though given in the majority of cases, did not add anything.

Concerning the localities "Arusha" and "Kilimanjaro", it must be supposed that these are general areas rather than exact collecting localities. Also the two Guso Nyiro series, from the Smithsonian Institution, are labelled respectively "Northern Guso Nyiro, Sotik" and "Southern Guso Nyiro, Sotik". This is confusing, as the town of Sotik lies considerably west of the Southern Guso Nyiro, and is nowhere near the Northern Guso Nyiro. Both indications must remain uncertain for the time being.

Applying the "75% rule"¹, one finds that the Kiboko, Tsavo and Tana populations are indistinguishable, and that the Karagwe and Serengeti groups are also indistinguishable, but perfectly distinguishable from the first three. We thus have two groups in East Africa, a small and a large one, with the Eyasi, Arusha and Kilimanjaro groups being intermediate. The Baringo skulls are beyond the limits of variability of either of the two groups, and the Lakimunda and Aberdare series are intermediate between Baringo and the Tana group. Turning now to populations outside the East African area, we find that, rather surprisingly, the S.E. African skulls (Natal and Cattedra, indistinguishable on ZUKOWSKY'S figures) are not significantly different from the large East African type (Karagwe and Serengeti). The Zambesi skulls show a higher variability than those from other areas; and this is indeed a somewhat heterogeneous category, formed from a series of skulls with no better locality than "Mashonaland" or "Mozambique". The West African skulls are very close to the small East African type, while both the Cape and Chobe skulls are perfectly distinct from all others.




Fig. 1. *Diceros bicornis* localities in East Africa, referred to in text. Not listed on map: Chertansang Hill; Archers Post; Longvya water. These localities are a little further north than the limits of the map, the first being near the Uganda border, the other two approximately due north of Mt. Kenya.

¹ See MAYR, LINSLEY & SINGER, 1953, Methods & Principles of Systematic Zoology, p. 146.

Clearly, we have here the basis of a taxonomic grouping. A single widespread race is indicated, distributed from Transkei in the south to Lake Victoria in the north, with a very large race at the Cape, another rather large one on the Chobe river, two smaller ones respectively in East Africa to the east of the Rift Valley and in West Africa, and a large broad-skulled one to the northwest of the Kenya distribution.

Turning now to ZUKOWSKY'S table, one must look for evidence that the other "races" fall into one of these major types, or else form a further type. They may be dealt with one by one.

keitola. — The single skull attributed to this race is smaller than most of the Cape skulls but considerably larger than the so-called *D. b. keitola* var. *minor*. Its toothrow length, however, comes nearer the latter. Unless it is merely a small-toothed variant of the Cape form, it would seem to make a good intermediate between the Cape and the large Eastern and South-eastern type.

occidentalis. — Apart from a very large zygomatic breadth, the single skull of this form would belong easily with the Eastern and South-eastern type.

angolensis. — These skulls show no difference from the Eastern and South-eastern race. This makes it all the more likely that the broad zygomatic of the geographically intervening *occidentalis* are attributable to individual variation.

rossmuae. — This is one of the invalid "Eventualnamen" pointed out by MERRENS. In addition to the type in Berlin, there is also a skull labelled "Rovuma river" in the British Museum. Neither of the two shows any difference from the Eastern and South-eastern race.

nichabli. — The Holotype of this subspecies is from "between Engaruka and Serengeti", i. e. in the intergrade zone between the small and large East African races. The type itself, however, agrees on a purely typological basis with the small race.

rendilis. — The two Northern Guano Nyiro specimens agree better with the small East African race, while the Jubaland skull which is attributed to *rendilis* is of the large type.

lidoensis. — The Holotype, from Shanbe, and a skull in the Smithsonian Institution from Gondokoro, both are strikingly reminiscent, in their broad zygomatic and occiput, of the Baringo skulls in Kenya.

somalensis. — The single skull attributed to this race does not fit clearly into any of the other forms, being especially unique in its very narrow zygomatic breadth. Two subadult skulls in Vienna, from Mersi in Ogaden, agree well with the Somali skull, being (size for size) only a little broader at the zygomatic.

brucii and *atbarensis*. — These two races must be re-shuffled. The two skulls listed as *atbarensis* are both decidedly subadult: on the other hand some of the skulls listed under *brucii* from "Abyssinia" have a more exact locality. Number 74, 11. 2. 2. is from Gachir Serit, and thus a virtual topotype of *brucii*; while no. 69, 2. 2. 14 is from Bejki, Anseba valley, no. 76, 9, 26, 6 is from Athbara river, and no. 79, 11, 29, 4 from Bogos. A further skull from Bogos is in the Cambridge museum. All these last would belong to *atbarensis* rather than to *brucii*, and they are indeed — those that are fully adult — a little smaller, but not very markedly, and a size variation covering both supposed races can be found within either of the East African forms. In general, the skulls from Ethiopia are not unlike the Somali and Ogaden skulls, but slightly broader.

Before compiling synonyms and calculating subspecies parameters, one must look at the photographs of rhinoceroses in the flesh, to determine whether any external features serve to characterise one race or another. It is felt by the present author that in characterising the subspecies externally, ZUKOWSKY did not sufficiently take age factors into account, and that such things as depth of skin-folds and costal grooving would be very dependant upon age. None the less three habitus-types do appear distinguishable:

1. The general East, South-east and South-west African type, with a short, compact body, well-marked skin-folds and large head.
 2. The *brucii-atbarensis* type, also the photograph identified as *rendilis* on page 120, with a much longer, more slender body, small head and comparative lack of skin-folds.
 3. The West-African type, *D. b. longipes*, with its long legs, smooth skin, short body and characteristic thickened base to the anterior horn.
- It may be noted that the external appearance of the Cape and Chobe races is unknown.
- The subspecies of *Diceros bicornis* may now be allotted and defined. In the calculation of the parameters (Table 2), all single-specimen localities were allotted on the basis of geography, and it was found that, e. g. the skulls from Taveta and Amboseli were morphologically also of the Kiboko-Tsavo-Tana type. In some cases, however, where a delimitation of range was involved, skulls had to be allotted typologically, e. g. the large size of the skull from Umba steppe was taken to mean that the large East African race encircles the small one to the south.

The subspecies may now be listed as follows, with full synonymy:

1. *Diceros bicornis bicornis* (Linnaeus, 1758). Cape black rhinoceros.
- 1758: *Rhinoceros bicornis* Linnaeus, Syst. Nat. ed. 10, 1:56. "India" (= Cape of Good Hope according to HOWAS).
- 1803: *Rhinoceros africanus* Blumenbach, Man. H. N. 1:156. Cape of Good Hope.

Table 2

Subspecies of *Diceros bicornis*: Means and Standard deviations

Measurements of intergrade specimens and populations are given in Table 1

	Greatest length	Basal length	Zygomatic breadth	Toothrow length	Occiput breadth
1. Adult: third molars in <i>aequalis</i> suture closed					
<i>bicornis</i> (5)	667.0 ± 37.7	626.3 ± 25.3	372.5 ± 13.4	315.3 ± 17.6	218.8 ± 23.6
<i>diobienensis</i> (4)	615.0 ± 15.6	565.0 ± 8.4	352.0 ± 9.9	282.3 ± 7.4	197.5 ± 6.3
<i>minor</i> (23)	576.0 ± 17.0	545.5 ± 16.9	330.4 ± 10.5	254.1 ± 13.0	188.2 ± 12.1
<i>nichabli</i> (22)	552.6 ± 20.9	514.2 ± 14.2	326.8 ± 9.8	251.2 ± 13.1	186.2 ± 10.1
<i>lidoensis</i> (6)	597.8 ± 20.9	557.5 ± 14.2	347.7 ± 11.2	253.7 ± 13.9	210.2 ± 6.3
<i>brucii</i> (10)	570.3 ± 18.9	531.8 ± 13.7	315.4 ± 9.3	251.4 ± 13.6	182.5 ± 10.6
<i>longipes</i> (4)	541.3 ± 17.5	522.7 ± 10.8	324.3 ± 3.8	257.5 ± 3.5	191.0 ± 5.5
2. Juvenile: third molars in <i>aequalis</i> suture open					
<i>bicornis</i> (2)	577.5	543.0	335.0		196.0
<i>minor</i> (2)	541.5	521.3	313.5		179.0
<i>nichabli</i> (1)	519	503	316		170
<i>lidoensis</i> (2)	537.5	516.5	324.0		182
<i>brucii</i> (1)	523	502	292		166
3. Late juvenile: third molars unerupted					
<i>minor</i> (3)	489.4	503.4	291.8		167.8
<i>nichabli</i> (2)	476.0	465.0	298.0		166
<i>lidoensis</i> (2)	501.0	503.5	320.0		—
<i>brucii</i> (4)	489.3	462.3	279.8		164.5
<i>longipes</i> (3)	490.0	492.3	294.3		169.7




Fig. 2. Distribution of subspecies of *Diceros bicornis*.

1836: *Rhinoceros keitloa* A. Smith, Rep. Exped. Expl. C. Africa, 44, "Country north and south of Kurriehaine" (Marico district, W. Transvaal). Although probably an intergrade between this race and *D. b. minor*, this name may be fixed as a synonym of the present race.

- 1837: *Rhinoceros keitloa* A. Smith, Cat. S. Afr. Mus. 7, 180 miles N. E. of Latrakoo (Kuruman).
- 1842: *Rhinoceros bicornis* var. B. *Rhinoceros gordonii* Lesson, Nouv. Tabl. Règne Anim. Mamm., 159. Based on BLAINVILLE's "Rhinoceros de Gordon", from environs of the Cape.
- 1845: *Rhinoceros niger* Schinz, Synops. Mamm., 2:335. Chantop (see ZUKOWSKY, p. 50). The specimen quoted by ZUKOWSKY, from near the type locality, has a mandibular length of 520 mm., as big as any Cape specimen.
- 1845: *Rhinoceros campeni* Schinz, loc. cit. Cape of Good Hope.
- 1898: *Rhinoceros bicornis capensis* Trouessart, Cat. Mamm. Viv. Foss. 757. Cape.
- Distribution: Fish River, Cape of Good Hope, Hottentots River (Beaufort West; an incomplete "pick-up" skull in British Museum), to middle latitudes in Southwest Africa.
2. *Diceros bicornis doliensis* Zukowsky, 1964.
- 1964: *Diceros bicornis doliensis* Zukowsky, Zool. Garten (N. F.) 30:79. Konsum-bia, parent-streams of the Loma, tributary of the Cuando, Angola.
3. *Diceros bicornis minor* (Drummond, 1876). Large East and South-East African rhinoceros.
- 1876: *Rhinoceros bicornis major* & *minor* Drummond, P. Z. S. 109. *R. b. minor* ("all country south-east of Zambesi", Zululand selected by Zukowsky) has priority because of selection by Zukowsky as first reviser. *R. b. major*, from Black Umfolosi to Limpopo.
- 1876: *Rhinoceros keitloa* Drummond, loc. cit. Black Umfolosi to Limpopo.
- 1893: *Rhinoceros holmwoodi* Sclater, P. Z. S. 517. Udalla, 50 miles S. of Speke Gulf.
- 1922: *Opisiceros occidentalis* Zukowsky, Arch. Naturgesch. 88 A, 7:162. Kaokoveld-Kunene region.
- 1947: *Diceros bicornis punyama* Potter, Field, 190:385. Hluhluwe Game Reserve.
- 1964: *Diceros bicornis angolensis* Zukowsky, Zool. Garten 30:73. Virui watehole, Mossamedes, Angola.
- 1964: *Diceros bicornis nyasae* Zukowsky, loc. cit. 93. North end of L. Nyasa, Karonga, and Niam-niam land, east of L. Rukwa. Invalid; proposed by implication.
- 1964: *Diceros bicornis yocennae* Zukowsky, loc. cit. 94. Inland from Mkhindani. Invalid; proposed by implication.
- Note. The theory that two forms of Black Rhinoceros coexist in Southeast Africa is discussed by ZUKOWSKY (pp. 41-5), who recognises a large *D. b. keitloa* (synonym *major* Drummond) and a small *D. b. keitloa* var. *minor* (synonym *punyama* Potter). If two forms do exist side by side, they cannot be conspecific. However Miss ADRIENNE ZHILMAN was kind enough to make enquiries while in South Africa on my behalf; game wardens, who knew the relevant literature, were unanimous in agreeing that only one form existed. Perhaps there was confusion with pairing of subadult animals; FOSTER (1965) indicates that the probable breeding population starts with animals with third molars only slightly worn, i. e. not full-sized. Accordingly, all three of DRUMMOND's names will refer to this race: which ranges from Transkei and northern S. W. Africa to L. Victoria.
4. *Diceros bicornis michaelsi* Zukowsky, 1964. Small East African black rhinoceros.
- 1964: *Diceros bicornis michaelsi* Zukowsky, Zool. Garten 30:115. Between Engaraka and Serengeti. Although the type locality is in the intergrade zone between

this race and the last, the skull is typical for the small race; and it is felt apt to continue to honour MICHAEL GRZIMEK, for whom the race is named.

1964: *Diceros bicornis renaldi* Zukowsky, loc. cit. 122. Northern Guaso Nyiro.

Distribution: Kenya, east of the Rift valley, into extreme northern Tanzania.

5. *Diceros bicornis brucei* (Lesson, 1842). Somali black rhinoceros.

1842: *Rhinoceros bicornis* var. *A. R. brucei* Lesson, Nouv. Tabl. Règne Anim. Mamm. 159. Based on Blainville's "Rhinoceros de Bruce" from Tcherkin, between Bahr Saliam and Abbara rivers.

1870: *Rhinoceros kerilwa* Blanford, Geol. Zool. Abyssinia, 243. Nor. of A. Smith, 1836.

1897: *Rhinoceros bicornis somaliensis* Potocki, Notarki Mysiowski z Atryki, 82. Somaliland.

1947: *Diceros bicornis palustris* Benzon, Field, 189:329. Near Aweng, north of Lol river, Bahr-el-Ghazal. Not mentioned by Zukowsky. Measurements fit well with present race.

1964: *Diceros bicornis atbarensis* Zukowsky, Zool. Garton 30:141. Anseba valley, Eritrea.

Distribution: northern Somalia and Ogaden to Sennaar and Bahr-el-Ghazal district.

6. *Diceros bicornis ladaensis* Zukowsky, 1964:²

1914: *Diceros bicornis somaliensis* J. Allen, Bull. Amer. Mus. N. H. 33:340. Cherangani Hills, Kenya. Not of POTOCKI, 1897.

1964: *Diceros bicornis ladaensis* Zukowsky, Zool. Garton, 30:124. Shamba, upper Nile, near Lado. Proposed by implication; invalid (MERREVS, 1966).

Diagnosis: a large race, larger than *minor* but smaller than *dobienisi*, with very broad zygomatic arches and occiput. See Table 2.

Distribution: Shamba and Gondokoro to Baringo district, thence to L. Naivasha.

7. *Diceros bicornis longipes* Zukowsky, 1949.

1949: *Diceros bicornis longipes* Zukowsky, Arche Noah, 1:16. Mlogrum (Chad Republic).

It must be admitted that there is considerable uncertainty in trying to fit clines into a taxonomic scheme. The three-way clinal variation in East Africa between *minor*, *michaeli* and *ladaensis* does, nonetheless, satisfactorily permit the breaking-up of the specimens into three groups — subspecies as above — and their intergrades. Fig. 1 breaks down taxonomically as follows:

- D. b. minor*: A, B, C, D, F, L.
- minor* — *michaeli* intergrades: E, G, H, I, K.
- D. b. michaeli*: J, M, N, O, P, Q, R, S, T.
- michaeli* — *ladaensis* intergrades: U, V.
- D. b. ladaensis*: W, X, Y, Z.

In addition to those shown on the map, Archers Post and Longava water are localities for subspecies *michaeli* and Cherangangi Hills (and Lado and Gondokoro) for *ladaensis*. A skull labelled "Nairobi" — probably a collecting centre rather than

² The name *Diceros bicornis somaliensis* J. Allen, 1914, is of course much earlier than *ladaensis* Zukowsky, 1964, but it is preoccupied by *somaliensis* Potocki, 1897 — synonym of *D. b. brucei* according to me, so that *ladaensis* must remain as the valid name.

an actual locality — is also typical for *ladaensis*. Finally, skulls from Jubaland and from Lamu in Ogaden are intergrades between *michaeli* and *brucei*.

Information on the index-cards of the Powell-Cotton collection gives the shoulder-height of a rhinoceros from East of Lake Baringo as 150 cm. If this is taken by a comparable technique as the measurements for various races given by Zukowsky, this would indicate a small bodily size for a large skulled animal. From Zukowsky's data it would seem that shoulder heights for other races are as follows:

<i>D. b. bicornis</i>	202–224 cm (2)
<i>D. b. dobienisi</i>	170–175 cm (2)
<i>D. b. minor</i>	157–179 cm (16)
<i>D. b. michaeli</i>	132–155 cm (7)
<i>D. b. brucei</i>	137–142 cm (2)
<i>D. b. longipes</i>	146–147 cm (2) (Powell-Cotton).

Summary

The classification of the subspecies of the Black Rhinoceros, *Diceros bicornis*, is reviewed in the light of a re-interpretation of ZUKOWSKY'S (1964) data, with additional data collected by the author. It is concluded that in East Africa there is a size-increase cline from east to west in southern Kenya and northern Tanzania, and another size-increase cline together with marked increase in skull breadth from south-eastern to north-western Kenya. The three extremes are given subspecific rank: one of the three is indistinguishable from specimens in Zambia, Malawi, and Zululand and Angola. There are two additional subspecies in southern Africa, recognizable by skull characters; while two others, not so easily distinguishable by their skulls but easily characterised externally, inhabit respectively the Somalia-north Sudan region and the Lake Chad area. Synonymy and skull measurements are given; one of ZUKOWSKY'S invalid names, *Diceros bicornis ladaensis*, is here used for the first time as the name of a definite race, and so obtains validity.

Zusammenfassung

Die Klassifikation der Unterarten des Schwarzen Nashorns, *Diceros bicornis*, wird nach den Daten von ZUKOWSKY (1964) und zusätzlich, vom jetzigen Autor gesammelten neu interpretiert. Es ergibt sich, daß in Ostafrika die Größe von Osten nach Westen zunimmt vom südlichen Kenya zum nördlichen Tanzania, und daß sich eine weitere Größenzunahme findet — gleichzeitig mit merklicher Breitenzunahme des Schädels — vom südöstlichen nach dem nordwestlichen Kenya. Den drei Extremlinien wird subspezifischer Rang zugesprochen. Eine dieser drei Formen ist nicht zu unterscheiden von Exemplaren aus Zambia, Malawiland, Zululand und Angola. Zwei weitere Subspezies im südlichen Afrika sind durch Schädelmerkmale zu diagnostizieren. Zwei andere, die nach dem Schädel nur schwer, aber nach der äußeren Erscheinung leicht zu unterscheiden sind, bewohnen resp. die Somali-Nord Sudan-Region und das Tschadsee-Gebiet. Synonyme und Schädelmessungen werden gegeben. Einer von ZUKOWSKY'S invaliden Namen, nämlich *Diceros bicornis ladaensis*, wird hier zum ersten Mal als Name einer definierbaren Rasse benutzt und wird dadurch valide.

Acknowledgements

The author is greatly indebted to Miss J. H. KINE of the Osteology Dept., British Museum (Natural History); Mr. R. D. NOKSAN of the Cambridge University Museum, Zoology; Prof. A. E. W. MILES of the Odontology Museum, Royal College of Surgeons; Mr. L. R. BARKER of the Powell-Cotton Museum, Bretherton, Kent, England; Dr. Th. HALTENSORH and Herr E. THUMER of the Zoologische Staatssammlung, Munich; Dr. KURT BAUER of the Naturhistorisches Museum, Vienna; Dr. E. W. BEAUFREY of the Universitet, Zoologisk Museet, Copenhagen; Dr. R. VAN GELDER and Dr. SIDNEY E. ANDERSON of the Mammalogy Department, American Museum of Natural History, New York; and Dr. C. O. HANDLEY of the Mammal Division, Smithsonian Institution, Washington, D. C.